arXiv:1703.02963v1 [math.PR] 8 Mar 2017

Central Limit Theorem for a Self-Repelling Diffusion

Carl-Erik Gauthier*
Department of Mathematics
University of Washington, USA

March 9, 2017

Abstract

We prove a Central Limit Theorem for the finite dimensional distributions of the displacement for the 1D self-repelling diffusion which solves

$$dX_t = dB_t - \big(G'(X_t) + \int_0^t F'(X_t - X_s)ds\big)dt,$$

where B is a real valued standard Brownian motion and $F(x) = \sum_{k=1}^n a_k \cos(kx)$ with $n < \infty$ and $a_1, \cdots, a_n > 0$.

In dimension $d \geq 3$, such a result has already been established by Horváth, Tóth and Vetö in [3] in 2012 but not for $d = 1, 2$. Under an integrability condition, Tarrès, Tóth and Valkó conjectured in [6] that a Central Limit Theorem result should also hold in dimension $d = 1$.

Keywords: Central Limit Theorem, Self-Repelling Diffusion
MSC2010: 60F05, 60K35, 60H10, 60J60

1 Introduction

In this short note, we aim to prove the Central Limit Theorem (denoted by CLT in the sequel) for the finite dimensional distribution of the displacement for the 1D self-repelling diffusion solving

$$dX_t = dB_t - \big(G'(X_t) + \int_0^t F'(X_t - X_s)ds\big)dt, \quad X_0 = 0, \tag{1}$$

where B is a real valued standard Brownian motion, $G(x) = \sum_{k=1}^n \big(u_k \cos(kx) + v_k \sin(kx)\big)$ and $F(x) = \sum_{k=1}^n a_k \cos(kx)$ with $n < \infty$ and $a_1, \cdots, a_n > 0$.

*Contact email: carlgaut@uw.edu or carlerik.gauthier@gmail.com
On leave from Université de Neuchâtel, Switzerland

Roughly speaking, *self-repelling diffusions* (as considered here) are time continuous stochastic processes which solve an inhomogeneous stochastic differential equation whose drift part is evolving in time according to the whole past history of the process in such a way that it tends to push the diffusing particle away from the most visited sites.

Under the assumptions made on F and G, the Law of Large Number has already been established in [1, Theorem 2 and Remark 1], namely

$$\lim_{t\to\infty} \frac{X_t}{t} = 0 \ a.s.$$

A question that one may then ask is whether or not a CLT result holds. The purpose of this note is to provide a positive answer to it.

To the author's knowledge it is the first time that such a result is obtained for a 1D self-repelling diffusion, despite being conjectured in 2012 by Tarrès, Tóth and Valkó ([6, Theorem 2 and its remark]) under a positivity[1] and an integrability[2] condition. Nevertheless, Horváth, Tóth and Vetö were able to prove in 2012 a CLT in dimensions $d \geq 3$ ([3, Theorem 2]) by using the Kipnis-Varadhan's CLT result for additive functionals via a *graded sector condition* which turns out to fail in dimension 1.

Before turning to the presentation of the result, let us briefly make a link between the positivity condition from [6] and the positivity of the coefficients $a_1, \cdots, a_n$. Let $b \in L^1(\mathbb{R}) \cap C^\infty(\mathbb{R})$ be a function satisfying the positivity condition from [6] and let

$$\varphi_{2\pi}(b)(x) = \sum_{n=-\infty}^{\infty} b(x + 2\pi n)$$

be the $2\pi-$periodization transform of b. It is an exercise in Fourier analysis to show that

$$\varphi_{2\pi}(b)(x) = \frac{1}{2\pi} \sum_{k\in\mathbb{Z}} \hat{b}\big(\frac{k}{2\pi}\big) e^{ikx}.$$

Because b is even, we have

$$\varphi_{2\pi}(b)(x) = \frac{\hat{b}(0)}{2\pi} + \frac{1}{\pi} \sum_{k\geqslant 1} \hat{b}\big(\frac{k}{2\pi}\big) \cos(kx).$$

The paper is organised as follows. In Section 2, we state the CLT result and present the tools and concepts involved while its proof is presented in Section 3.

2 Tools, notation and results

In this section, we introduce the mathematical background that is necessary to present the main result.

[1] The Fourier transform of F is nonnegative

[2] $\rho^2 := \int_{-\infty}^{\infty} p^{-2} \hat{F}(p) dp < \infty$

Following the same idea as in [1], set $U_j(t) = u_j + \int_0^t \cos(jX_s)ds$ and $V_j(t) = v_j + \int_0^t \sin(jX_s)ds$. With these new variables, we obtain the following system

$$\begin{cases} dX_t = dB_t + \sum_{j=1}^n ja_j\Big(\sin(jX_t)U_j(t) - \cos(jX_t)V_j(t)\Big)dt \\ dU_j(t) = \cos(jX_t)dt,\ j = 1,\dots,n. \\ dV_j(t) = \sin(jX_t)dt,\ j = 1,\dots,n. \end{cases} \tag{2}$$

Since for all $j = 1\cdots,n$, the functions $x \mapsto \cos(jx)$ and $x \mapsto \sin(jx)$ are 2π-periodic, we can replace X_t by $\Theta_t = X_t \pmod{2\pi} \in S^1$, where $S^1 = \mathbb{R}/2\pi\mathbb{Z}$ denotes the 1-dimensional flat torus. This replacement allows us to use the framework from [1].

In order to shorten the notation, we let U_t and V_t denote the vectors

$$U_t = \big(U_1(t),\cdots,U_n(t)\big) \qquad \text{and} \qquad V_t = \big(V_1(t),\cdots,V_n(t)\big).$$

Summarizing the main results from [1], we have

Theorem 1 (Theorem 5 and Theorem 6, [1])**.** *) Let $(P_t)_{t\geqslant 0}$ be the semi-group associated to the process $\Big((\Theta_t,U_t,V_t)\Big)_{t\geq 0}$ and $P_t((\theta_0,u_0,v_0),d\theta dudv)$ its transition probability. Then*

1) The unique invariant probability measure is

$$\mu(d\theta dudv) = \nu(d\theta)\otimes\frac{e^{-\Phi(u,v)}}{C}dudv,$$

where $\Phi(u,v) = \frac{1}{2}\sum_{k=1}^n a_k k^2(u_k^2+v_k^2)$, C is a normalization constant and $\nu(d\theta)$ is the uniform probability measure on $S^1 = \mathbb{R}/2\pi\mathbb{Z}$.

2) Let $\mu_t = \mathcal{L}(\Theta_t,U_t,V_t)$ denote the law of (Θ_t,U_t,V_t). Then for any initial distribution μ_0, μ_t converges to μ in total variation.

3) For every $\eta > 0$ and $g \in L^2(\mu)$

$$\|P_t g - \int g(\theta,u,v)\mu(d\theta dudv)\|_{L^2(\mu)} \leqslant \sqrt{1+2\eta}\|g - \int g(\theta,u,v)\mu(d\theta dudv)\|_{L^2(\mu)}e^{-\lambda t},$$

where

$$\lambda = \frac{\eta}{1+\eta}\frac{K_1}{1+K_2+K_3},$$

with explicit constants K_1, K_2 and K_3.

In this paper, we will adopt the same point of view as in [6]: *the environment seen from the particle*. For that purpose, introduce the following new variables

$$C_j(t) = U_j(t)\cos(jX_t) + V_j(t)\sin(jX_t) = \left\langle \begin{pmatrix} U_j(t) \\ V_j(t)\end{pmatrix}, \begin{pmatrix}\cos(jX_t)\\ \sin(jX_t)\end{pmatrix}\right\rangle \tag{3}$$

and

$$S_j(t) = \sin(jX_t)U_j(t) - \cos(jX_t)V_j(t) = \Big\langle \begin{pmatrix} U_j(t) \\ V_j(t) \end{pmatrix}, \begin{pmatrix} \sin(jX_t) \\ -\cos(jX_t) \end{pmatrix} \Big\rangle. \tag{4}$$

So, if we denote by η_t the potential viewed from the particle's position, i.e

$$\eta_t(x) = F_t(x + X_t) = \int_0^t \sum_{k=1}^n a_k \cos(k(x + X_t - X_s))ds + G(x + X_t),$$

then

$$\eta_t(x) = \sum_{k=1}^n a_k \Big(C_k(t)\cos(kx) - S_k(t)\sin(kx) \Big).$$

Moreover, this allows us to rewrite X_t as

$$X_t = B_t + \int_0^t \sum_{k=1}^n k a_k S_k(u)du = B_t + \int_0^t \varphi(\eta'_u)du,$$

where $\varphi : \Omega \to \mathbb{R}$ is defined by $\varphi(\omega) = \omega(0)$ and Ω is the vector space spanned by the functions $\cos(kx)$ and $\sin(kx)$ for $k = 0, 1, \cdots n$.

Before turning to the results, let us introduce the following notation. We denote by T_t the semigroup induced by the process

$$((C_t, S_t))_{t\geqslant 0} := \Big(\big(C_1(t), S_1(t), \cdots, C_n(t), S_n(t) \big) \Big)_{t\geqslant 0}$$

and by G its infinitesimal generator. For an operator R, we denote its domain by $D(R)$.

Given a probability measure π over $\mathbb{R}^{2n}$, we denote by $L^2(\pi)$ the space $L^2(\mathbb{R}^{2n}, \pi)$, by $\Big\langle ., . \Big\rangle_{L^2(\pi)}$ the associated inner product and by $\Big\| . \Big\|_{L^2(\pi)}$ the induced L^2-norm. Finally, we denote by $\mathbb{P}_\pi$ the law of the process with initial distribution $\pi = \mathcal{L}\big((C_0, S_0)\big)$ and by $\Longrightarrow$ the convergence in distribution.

Proposition 1.

1. *For any smooth function f having compact support, we have*

$$\begin{aligned} Gf(c,s) \;=\; & \frac{1}{2}\sum_{j=1}^n j^2 \big(s_j^2 \partial_{c_j c_j} f + c_j^2 \partial_{s_j s_j} f \big) - \frac{1}{2}\sum_{k\neq j} jk \big(s_j s_k \partial_{c_k c_j} f + c_j c_k \partial_{s_j s_k} f \big) \\ & - \sum_{j,k=1}^n jk s_j c_k \partial_{c_j s_k} f + \Big(\sum_{k=1}^n k a_k s_k \Big) \sum_{j=1}^n j \big(- s_j \partial_{c_j} f + c_j \partial_{s_j} f \big) + \sum_{j=1}^n \partial_{c_j} f. \end{aligned}$$

2. *The process $((C_t, S_t))_{t\geqslant 0}$ admits a unique invariant probability measure of the form*

$$\pi(dcds) = \frac{e^{-\Phi(c,s)}}{C} dcds$$

where $\Phi(c,s) = \frac{1}{2}\sum_{k=1}^n a_k k^2 (c_k^2 + s_k^2)$ and C is the normalizing constant.

3. For any function $f \in L^2(\pi)$, we have

$$\Big\| T_t f - \int_{\mathbb{R}^{2n}} f(c,s)\pi(dcds) \Big\|_{L^2(\pi)} \leq \sqrt{3} \Big\| f - \int_{\mathbb{R}^{2n}} f(c,s)\pi(dcds) \Big\|_{L^2(\pi)} e^{-\lambda t},$$

where $\lambda = \frac{1}{2}\frac{K_1}{1+K_2+K_3}$ and the constants K_1, K_2 and K_3 are those from Theorem 1.

Proof.

1. It follows from Itô's formula and [5, Propositions VII.1.6 and VII.1.7].

2. The fact that $\pi(dcds)$ is an invariant probability measure follows from Theorem 1 as well as from the equations (3) and (4). Indeed, for any $A_j \in \mathcal{B}(\mathbb{R}^2)$, we have by rotation invariance of the Gaussian measure

$$\begin{aligned} \pi\Big(A_1 \times \cdots \times A_n\Big) &= \mathbb{P}\Big(\big(C_j(t), S_j(t)\big) \in A_j \, \forall j = 1, \cdots, n\Big) \\ &= \mathbb{P}\Big(\big(U_j(t), V_j(t)\big) \in A_j \, \forall j = 1, \cdots, n\Big) \\ &= \mathbb{P}\Big(\Theta_t \in \mathbb{S}^1, \ \big(U_j(t), V_j(t)\big) \in A_j \, \forall j = 1, \cdots, n\Big) \\ &= \mu\Big(\mathbb{S}^1 \times A_1 \times \cdots \times A_n\Big) \end{aligned}$$

   Concerning the uniqueness, let ν be an invariant probability measure for the process $((C_t, S_t))_{t \geqslant 0}$. Then define on $S^1 \times \mathbb{R}^n \times \mathbb{R}^n$ the probability measure $\mu_0(d\theta dudv) = \delta_0 \otimes \nu(dudv)$ and sample (Θ_0, U_0, V_0) according to μ_0.

   By Theorem 1, μ_t converges to μ in total variation. In particular the marginal law of μ_t corresponding to (U_t, V_t) converges to π. Thus $\nu = \pi$.

3. Let $f : \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}$ and define a function $g : S^1 \times \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}$ by

$$g(\theta, u, v) = f(c, s),$$

   where the pairs (c_j, s_j) are defined as in (3) and (4). Applying Itô's formula to (3) and (4) yields

$$d\begin{pmatrix} C_1(t) \\ S_1(t) \\ C_2(t) \\ S_2(t) \\ \vdots \\ C_n(t) \\ S_n(t) \end{pmatrix} = \begin{pmatrix} -S_1(t) \\ C_1(t) \\ -2S_2(t) \\ 2C_2(t) \\ \vdots \\ -nS_n(t) \\ nC_n(t) \end{pmatrix} \Big(dB_t + (\sum_{k=1}^{n} k a_k S_k(t))dt\Big) - \begin{pmatrix} C_1(t) \\ S_1(t) \\ 4C_2(t) \\ 4S_2(t) \\ \vdots \\ n^2 C_n(t) \\ n^2 S_n(t) \end{pmatrix} dt + \begin{pmatrix} 1 \\ 0 \\ 1 \\ 0 \\ \vdots \\ 1 \\ 0 \end{pmatrix} dt.$$

Thus the evolution of (C_t, S_t) does not depend on the dynamic of Θ_t. Therefore

$$\begin{aligned} T_t f(c,s) &= \mathbb{E}\Big(f(C_t,S_t) \mid C_0 = c,\, S_0 = s \Big) \\ &= \mathbb{E}\Big(f(C_t,S_t) \mid \Theta_0 = \theta,\, C_0 = c,\, S_0 = s \Big) \\ &= \mathbb{E}\Big(f(C_t,S_t) \mid \Theta_0 = \theta,\, U_0 = u,\, V_0 = v \Big) \\ &= \mathbb{E}\Big(g(\Theta_t,U_t,V_t) \mid \Theta_0 = \theta,\, U_0 = u,\, V_0 = v \Big) \\ &= P_t g(\theta,u,v), \end{aligned}$$

where the pairs (u_k, v_k) are such that $c_k = u_k \cos(k\theta) + v_k \sin(k\theta)$ and $s_k = u_k \sin(k\theta) - v_k \cos(k\theta)$.

The statement follows then from Theorem 1 with $\eta = 1$.

□

Our CLT result is the following

Theorem 2.

1. *$\sigma^2 := \lim_{t\to\infty} \frac{\mathbb{E}_\pi(X_t^2)}{t}$ exists and it satisfies*

$$1 \le \sigma^2 \le 1 + 2\Big(\sum_{j=1}^{n} \frac{a_j}{j^2} \Big). \tag{5}$$

2. *For any $0 < t_1 < \cdots < t_n < \infty$, we have*

$$\Big(\frac{X_{t_1/\varepsilon}}{\sigma^2 t_1/\varepsilon}, \cdots, \frac{X_{t_n/\varepsilon}}{\sigma^2 t_n/\varepsilon} \Big) \overset{\varepsilon\to 0}{\Longrightarrow} \Big(W_{t_1}, \cdots, W_{t_n} \Big) \tag{6}$$

under $\mathbb{P}_\pi$, where W is a real valued standard Brownian motion.

3 Proof of Theorem 2

Throughout this section, we let $g, h : \mathbb{R}^n \to \mathbb{R}$ denote the functions defined by

$$g(c,s) = \sum_{k=1}^{n} k a_k s_k \qquad \text{and} \qquad h(c,s) = \sum_{k=1}^{n} a_k c_k.$$

3.1 Proof of Part 1

First of all, by repeating the arguments of [6], we have as at the beginning of [6, Section 4]

$$\begin{aligned}\mathbb{E}_\pi(X_t^2) &= t + \mathbb{E}_\pi\Big(\big(\int_0^t \sum_{k=1}^n k a_k S_k(u)du\big)^2\Big) \qquad (7)\\ &= t + 2\int_0^t (t-u)\mathbb{E}_\pi\Big(\big(\sum_{k=1}^n k a_k S_k(u)du\big)\big(\sum_{k=1}^n k a_k S_k(0)\big)\Big).\\ &= t + 2\int_0^t (t-u)\Big\langle T_u g, g\Big\rangle_{L^2(\pi)}. \qquad (8)\end{aligned}$$

By the Cauchy-Schwarz inequality and the third part of Proposition 1, $\Big\langle T_u g, g\Big\rangle_{L^2(\pi)}$ decreases exponentially fast to 0. Hence $\int_0^\infty \Big\langle T_u g, g\Big\rangle_{L^2(\pi)} du$ exists and, therefore, it yields

$$\lim_{t\to\infty} \frac{\mathbb{E}_\pi(X_t^2)}{t} = 1 + \int_0^\infty \Big\langle T_u g, g\Big\rangle_{L^2(\pi)} du := \sigma^2. \qquad (9)$$

Now that the existence of σ^2 is established, let us prove the bounds in (5). The lower bound is trivial since it follows from (7). In order to establish the upper bound, we follow the arguments presented in [4] based on the Kipnis-Varadhan's CLT theorem.

Letting G^* denote the adjoint of G in $L^2(\pi)$, $S = \frac{1}{2}(G + G^*)$ and $A = \frac{1}{2}(-G + G^*)$ denote the symmetric and the skew-symmetric part of G, we have for any smooth function f having compact support

$$\Big\langle Gf, f\Big\rangle_{L^2(\pi)} = \Big\langle Sf, f\Big\rangle_{L^2(\pi)} = -\frac{1}{2}\int \Big(\sum_{j=1}^n j(s_j\partial_{c_j} f - c_j\partial_{s_j} f)\Big)^2 d\pi. \qquad (10)$$

Hence, using the notation of [4], we have

$$\Big\langle Gf, f\Big\rangle_{L^2(\pi)} = -\|f\|_1^2. \qquad (11)$$

Because

$$\begin{aligned}\int h(c,s)\big(\sum_{j=1} j s_j\partial_{c_j} f - j c_j\partial_{s_j} f\big)d\pi &= \int \big(\sum_{j=1}^n j s_j\partial_{c_j} h - j c_j\partial_{s_j} h\big) f d\pi\\ &\quad - \int hf\big(\sum_{k=1} k^2 a_k c_k\big)d\pi\\ &\quad + \int hf\big(\sum_{k=1} k^2 a_k c_k\big)d\pi\\ &= \int gf d\pi \qquad (12)\end{aligned}$$

it follows from the Cauchy-Schwarz inequality that

$$\Big| \int g f d\pi \Big| \leq \|h\|_{L^2(\pi)} \|f\|_1. \tag{13}$$

Hence, with the notation of [4],

$$\|g\|_{-1} \leq \|h\|_{L^2(\pi)} = \sqrt{\sum_{j=1}^{n} \frac{a_j}{j^2}}. \tag{14}$$

Thus, the upper bound comes from Eq. (2.1.7) in [4].

3.2 Proof of Part 2

Set $\mathcal{S}_t = \int_0^t g(C_u, S_u)du$ and denote by $\mathfrak{s}_t^2$ its variance under π.

Since $\|T_t g\|_{L^2(\pi)}$ decreases exponentially fast to 0 by Proposition 1 and $\int g d\pi = 0$, then, by [2, Corollary 3.2.i], there exists a function $f \in D(G^{-1})$ such that $Gf = g$ and

$$\mathfrak{s}_t^2 = 2t\|f\|_1 = -\Big\langle g, f \Big\rangle_{L^2(\pi)}.$$

Because $\mathbb{E}_\pi\Big(S_t\Big) = 0$ for all $t \geq 0$, then $\|f\|_1 = 0$ implies that the process $t \mapsto \mathcal{S}_t$ is the null process $\mathbb{P}_\pi$-almost surely and the result is therefore trivial in that case.

If $\|f\|_1 > 0$, then by [2, Theorem 3.1], we have that for any $0 < t_1 < \cdots < t_n < \infty$

$$\Big(\frac{S_{t_1/\varepsilon}}{s_{t_1/\varepsilon}}, \cdots, \frac{S_{t_n/\varepsilon}}{s_{t_n/\varepsilon}}\Big) \overset{\varepsilon \to 0}{\Longrightarrow} \Big(W_{t_1}, \cdots, W_{t_n}\Big) \tag{15}$$

under $\mathbb{P}_\pi$, where W is a real valued standard Brownian motion.

Thus, the second part of Theorem 2 follows from the martingale approximation in the Kipnis-Varadhan approach as presented in [2].

Acknowledgement

I acknowledge financial support from the Swiss National Foundation for Research through the grants 200021_163072 and P2NEP2_171951.

I am grateful to Bálint Tóth for asking me wether or not a CLT result holds in the periodic case considered in this work.

References

[1] Benaïm M. and Gauthier C.E.: *Self-repelling diffusions on a Riemannian manifold*, Probab. Theory Relat. Fields., Article first published online: 06 May 2016, DOI: 10.1007/s00440-016-0717-1 (2016)

[2] Cattiaux, P., Chafaï, D., Guillin, A.: *Central Limit Theorems for Additive Functionals of Ergodic Markov Diffusions Processes*, ALEA, Vol 9(2), 337-382 (2012)

[3] Horváth, I., Tóth, B. and Vetö, B.: *Diffusive limits for "true" (or myopic) self-avoiding random walks and self-repellent Brownian polymers in $d \geq 3$*, Probab. Theory Relat. Fields., Vol. 153, Issue 3-4, 691-723 (2012)

[4] Olla S.:*Central Limit Theorems for tagged particles and for Diffusions in random environment*. In: Comets,F., Pardoux,E. (eds.) Milieux aléatoires Panor. Synthèses 12. Soc. Math. France, Paris (2001)

[5] Revuz, D., Yor, M.: *Continuous Martingales and Brownian Motion*, North-Holland Mathematical Library . Springer, 1999.

[6] Tarrès, P., Tóth, B., Valkó, B.: *Diffusivity bounds for 1d Brownian polymers*, Ann. Probab., vol. 40, Issue 2, 437-891 (2012)